

# Spectroscopy Analysis and Soil Pollution Study a round Balhaf Liquid Natural Gas, Yemen.

Khaled Laqal<sup>1,2</sup>, Siddig Tawaer Kafi<sup>3</sup>, Yousef. A. Alsabah<sup>4\*</sup>

<sup>1</sup> Department of Physics, faculty of Oil and Minerals, Shabwah University, Shabwah, Yemen

<sup>2</sup> Department of Physics, faculty of Science, Bahri University, Khartoum, Sudan

<sup>3</sup> Department of Physics, College of applied Science, Alneelain University, Khartoum, Sudan

<sup>4</sup> Department of Physics, Colleges Education and applied Science, Hajjah University, Hajjah, Yemen

**Abstract:** The aimed of study to determine the contaminants in soil resulting due to emissions that come from natural gas in the station of Balhaf which is it LNG project in Shabowah Gov, Republic of Yemen. Total 20 samples soil samples were collected at depth of approximately (0-10 cm) from different locations from the Balhaf natural gas station. These samples were analyzed by Laser Breakdown Plasma Spectroscopy (LIBS) to determine the samples toxic elements. The components of the toxic elements present in the soil samples were further confirmed by spectroscopic technique induced coupled plasma (ICP) and Fourier Transform Infrared spectroscopy (FTIR) techniques. The LIBS spectra showed the presence of different amounts of elements, including radioactive and toxic elements such as (Ac, Pu, Th, U) and (Al, Ba, Cd, Co, Cr, Cu, Hg, Mn, Ni, Pb, Tl, Zn), in all of the soil samples. The concentrations of the elements were calculated by ICP spectroscopy. The average concentrations of elements in the samples appeared to be 410 ppb for Al, 1.9 ppb for Ba, 2.6 ppb for Cd, 2.7 ppb for Co, 2.3 ppb for Cr, 257 ppb for Cu, 30 ppb for Mn, 6.8 ppb for Ni, 63 ppb for Th, 56 ppb for Tl and 6.1 ppb for Zn, but these values did not exceed normal levels for these elements in earth crust according to reference values such as ATSDR 2020, IAEA, 200. The presence of a high concentration of toxic elements in the soil around the natural gas facility, that indicates the presence of emissions from this facility that leads to the accumulation of these elements in the soil.

**Keywords:** LNG Project, LIBS, ICP, FTIR, Pollution, yemen.

## 1. INTRODUCTION

Contaminants are defined as inputs of alien and potentially toxic substances into the environment; not all contaminants cause pollution, as their concentrations may be too low. Pollutants are defined as anthropogenically introduced substances that have harmful effects on the environment [1]. Generally, environmental pollution is the contamination of air, water, or food in such a manner as to cause real or potential harm to human health or well-being, or damage or harm nonhuman nature without justification [2]. As pollutants interact with the

environment, they undergo physical and chemical changes and are ultimately incorporated into the environment. The environment acts as a continuum into which all waste materials are placed. The pollutants, in turn, obey the second law of thermodynamics: matter cannot be destroyed; it is merely converted from one form to [3]. Soil is a complex amalgam, a non-renewable natural resource because it cannot be re-created except within the context of geological timescales. It can be simply defined as the unconsolidated mineral or organic material on the immediate surface of the Earth that serves as a natural medium for the growth of land plants [1]. Heavy metals are highly hazardous to the environment and organisms. It can be enriched through the food chain. Once the soil suffers from heavy metal contamination, it is difficult to be remediated [4]. Lesley Fleischman, et al. in 2016 founds that (38) counties in 21 states from the United States (US) face a cancer risk that exceeds the U.S. Environmental Protection Agency EPA's one-in-a-million threshold level of concern [5-8]. Lisa M. McKenzie, et al. in 2012 found that Residents living  $\leq \frac{1}{2}$  mile from wells are at greater risk for health effects from natural gas development (NGD) than are residents living  $> \frac{1}{2}$  mile from wells. Sub-chronic exposures to air pollutants during well completion activities present the greatest potential for health effects. The subchronic non-cancer hazard index (HI) of 5 for residents  $\leq \frac{1}{2}$  mile from wells was driven primarily by exposure to trimethylbenzenes, xylenes, and aliphatic hydrocarbons. Chronic HIs were 1 and 0.4. for residents  $\leq \frac{1}{2}$  mile from wells and  $> \frac{1}{2}$  mile from wells, respectively [6-9] Most of the previous studies in the natural gas sector and its impact on environmental pollution were focused on air analysis upon estimates of pollutant emissions using meteorological and emission data [10][11]. Despite the rapid expansion in natural gas extraction and production, there are not studies on the impact of residual pollution in soil surrounding natural gas plant on the environmental and health impacts these processes might have. This paper was study of pollution from the effect of natural gas in the station of Balhaf, Yemen.

**2. RESEARCH METHOD**

**2.1 Study Area (Balhaf) and Soil Sampling**

Soils were sampled from the vicinity of the Liquid natural gas (LNG) project, which is located at Balhaf in the South of Yemen on the Gulf of Aden Balhaf at Shabwah governorate of Yemen (longitude 48.1802°E, latitude 13.9854°N). Balhaf is an industrial port town and an oasis in an area of coastal dunes in the Rudum Coastal Area of Yemen. It has palm trees and white sand, which gives way to fields of black lava and to the fishing port of Bir-Ali. Near the area of the Balhaf, there are several villages as well as the nomads, and practiced agriculture and sheep grazing, goats, and camels. The construction of the LNG facilities lasted from September 2005 to the end of 2009 [12]. The production started on 15 October 2009 [13][14]. However, the Yemen LNG plant has been shut down and stopped commercial production and exports since April 2015 because of the Yemeni civil war.

**2.2 Characterizations**

**A - Inductively coupled Plasma Technique (ICP)**

The inductively coupled Plasma Technique (ICP-AES) analyses of the 20 soil samples were carried out the laboratory of General Administration of Forensic Evidence (Khartoum, Sudan) using Shimadzu (ICPE-9000, Japan). In this investigation, the samples were dried at 105C for 3 h. About 0.2 g of sample was accurately weighed into a container made of PFA (a perfluoroalkoxy polymer), which was then placed in a microwave pressure vessel

**B - Laser Induced Breakdown Spectroscopy Technique (LIBS) Measurement**

LIBS was used to identify the heavy metals and radioactive elements which can exist in polluted soil. Soil samples were irradiated by a nitrogen laser, with an energy of 100 mJ, which is operating in the ultraviolet range (typically 337.1 nm). This technique is based on the concept that the laser can vaporize the surface of the material and exciting its atoms or molecules with the laser pulse. The excited energy that holds the atom at the higher energy level will be released and the atoms return to their ground state.

**C. Fourier Transform Infrared Technique (FTIR)**

The FTIR was used to identify the functional groups of soil samples. The spectra were recorded between 4,000 and 400 cm<sup>-1</sup> using FTIR (FTIR-8400S, Shimadzu) with 1 cm<sup>-1</sup> resolution. Pellets were prepared by mixing the soil samples with KBr. By applying sufficient pressure, the mixture is pressed into a transparent disc, the usual analyse / KBr ratio is ca. 1:100. FTIR spectra of soil samples were recorded.

**3. RESULTS AND DISCUSSIONS**

**3.1 Inductively coupled Plasma Technique (ICP-AES) Result.**

The soil samples were collected and analysed by ICP technique to determine the concentration of heavy metals (e.g., Al, Ba, Cd, Co, Cr, Cu, Mn, Ni, Th, Tl, and Zn). Aluminium (Al) levels, 410 ppb, were the highest compared with other elements as indicated in table (1). This amount has a lower value compared to the crustal average according to reference values such as ATSDR 1999, IAEA, 200 [15]. In addition, the results showed that the Barium (Ba) concentration is 1.9 ppb, which is lower than the crustal average [16]. Cadmium (Cd), Cobalt (Co), and chromium (Cr) are poorly concentrated in the all samples, the distribution ranging from 2.3 to 2.7 ppb. The concentration of these elements did not exceed the normal concentration in crustal. Copper concentration level was found at 57 ppb, thorium (Th) was 63 ppb, Manganese (Mn) was the concentration at 30 ppb and thallium (Tl) was 56 ppb the obtained values of these elements are well below than the normal concentration in crustal. The zinc (Zn) levels was 6.1 ppb and nickel (Ni) was 6.8 ppb, which are lower than the crustal abundance [17][18][19].

**Table (1).** Concentrations of elements in all samples by ICP

Element	Concentration ppb
Al	410
Ba	1.9
Cd	2.6
Co	2.7
Cr	2.3
Cu	57
Mn	30
Ni	6.8
Th	63
Tl	56
Zn	6.1

**3.2 Laser Induced Breakdown Spectroscopy Technique (LIBS) Result.**


for indication, after irradiation with 100 mJ pulse energy, the spectra were recorded in the region from 385 nm to 1055 nm. Atomic spectra database National Institute of standards and Technology (NIST) and Hand book of Basic Atomic Spectroscopic Data was used for the spectral analysis of the samples [20]. The obtained LIBS spectra covered wide range of the spectrum ranging from the ultraviolet (UV) region to the near-infrared (NIR) region. Figure (1) show the high intensity of elements in LIBS spectra at all samples. The spectra of the soil samples indicate the presence of different amounts of elements, including radioactive and heavy metals such as (Ac, Pu, Th, U) and (Al, Ba, Cd, Co, Cr, Cu, Hg, Mn, Ni, Tl, Zn). The

Actinium ions (Ac+1) appeared in one sample with low amount. Aluminium atoms appeared in some samples with low amount especially in the one sample. Barium ions (Ba+1) was detected in some in samples. Cadmium atoms (Cd) appeared in three samples only with non-considerable amount. Cobalt atoms (Co), has just appeared in three samples with a low amount, additionally cobalt ions (Co+1) appeared with the low amount in two samples, there are several different wavelengths for the cobalt appeared as shown in the. Chromium atoms were found at a different amount, and with different wavelengths. The copper ion (Cu+2) was found in all samples with a high amount compared with the previous elements. Mercury atoms (Hg) were found with the relatively little amount in fourteen samples, (this element is highly toxic even with little amount [21]). Manganese ion (Mn+1) was presented with the relatively little amount in five samples. The Nickel ion (Ni+3) was found in high amounts in all samples. Plutonium atoms (Pu) was found in very high amounts in all samples. Moreover, Thorium atoms (Th) were found at a different amount, not that, the wavelength (1049.282) nm appeared in all samples with relatively high amounts. Besides neutral atoms, ions of different amounts and ionization stages also were recorded. Thallium atoms (Tl) appeared in some samples with a low amount. Uranium atoms appeared in five samples with low amounts especially in sample 3, also uranium ions appeared with the low amounts in fourteen samples. The zinc atom (Zn) was found in a low amount in one sample only.

**3.2 Laser Induced Breakdown Spectroscopy Technique (LIBS) Result.**

for indication, after irradiation with 100 mJ pulse energy, the spectra were recorded in the region from 385 nm to 1055 nm. Atomic spectra database National Institute of standards and Technology (NIST) and Hand book of Basic Atomic Spectroscopic Data was used for the spectral analysis of the samples [20]. The obtained LIBS spectra covered wide range of the spectrum ranging from the ultraviolet (UV) region to the near-infrared (NIR) region. Figure (1) show the high intensity of elements in LIBS spectra at all samples. The spectra of the soil samples indicate the presence of different amounts of elements, including radioactive and heavy metals such as (Ac, Pu, Th, U) and (Al, Ba, Cd, Co, Cr, Cu, Hg, Mn, Ni, Tl, Zn). The Actinium ions (Ac+1) appeared in one sample with low amount. Aluminium atoms appeared in some samples with low amount especially in the one sample. Barium ions (Ba+1) was detected in some in samples. Cadmium atoms (Cd) appeared in three samples only with non-considerable amount. Cobalt atoms (Co), has just appeared in three samples with a low amount, additionally cobalt ions (Co+1) appeared with the low amount in two samples, there are several different

wavelengths for the cobalt appeared as shown in the. Chromium atoms were found at a different amount, and with different wavelengths. The copper ion (Cu+2) was found in all samples with a high amount compared with the previous elements. Mercury atoms (Hg) were found with the relatively little amount in fourteen samples, (this element is highly toxic even with little amount [21]). Manganese ion (Mn+1) was presented with the relatively little amount in five samples. The Nickel ion (Ni+3) was found in high amounts in all samples. Plutonium atoms (Pu) was found in very high amounts in all samples. Moreover, Thorium atoms (Th) were found at a different amount, not that, the wavelength (1049.282) nm appeared in all samples with relatively high amounts. Besides neutral atoms, ions of different amounts and ionization stages also were recorded. Thallium atoms (Tl) appeared in some samples with a low amount. Uranium atoms appeared in five samples with low amounts especially in sample 3, also uranium ions appeared with the low amounts in fourteen samples. The zinc atom (Zn) was found in a low amount in one sample only.



**Figure (1).** Examples of LIBS emission spectrum of first four samples.

**3.3 Fourier Transform Infrared Technique (FTIR) Result**

Figure (2) shows FTIR spectra of represented soil samples. The band due to the methyl functionalities vibrated at 875 cm-1 was used as a normalizing factor since no change is expected for the band of these groups. Several absorption peaks at different wave numbers were observed in the spectra of represented soil samples. The FTIR Spectrum contains the entire information about the molecular structure of the investigated samples. From the frequency and intensities of some spectral bands, it was possible to predict the kind of chemical groups present in the studied samples [22]. The results showed that the peaks around 2520 cm-1 and 1430 cm-1 are due to (O-H) (carboxylic acids) for most samples [23][24]. A normal range of carbonyl group (C=O) is 1800 to 1600

cm-1[25]. And the results showed that a strong transmittance band was at 1800 cm-1 within the range for all samples referred to as anhydrides [23]. Bands at 1625 cm-1, at 1430 cm-1, at 775cm-1 cm-1 and 710 cm-1 could be attributed to heterocyclic compounds (N-H), amines(NH2) and amides(N-H ) [26][23][27]. And appeared this results in all most samples as shown in the graphs from Figure (2), the band at 710 cm-1 corresponds to Ca-O bonds [26]. In addition, C-O group which has been observed in many compounds as Alcohol, ethers, esters, carboxylic acids, and anhydrides appeared at 1020 cm-1 with a strong band in all samples [23]. Also, the results of all samples indicated to the presence of aliphatic (C-N) stretching at 1020 cm-1 [27]. The results showed analysis group Nitrate NO2 asymmetric stretching at (710) cm-1 within the range (750–650) cm-1 in all samples except five samples [27]. The absorption bands at 875 cm-1 attributed to the out-of-plane bending of C-H, which observed in all samples [27][28]. In the fingerprint region, vibrations of many functional groups are observed, such as phosphorus compound at 460 cm-1 , and sulfur compounds at 520 cm-1, these groups appeared in all samples [27].







Figure (2). FTIR spectra of soil samples.

#### 4. Conclusion

This research focused on studies of soil pollution utilizing LIBS technology combined with ICP and FTIR. the search samples were soil from different inside of LNG project in Balhaf, coupled plasma technology showed the presence of the following elements (Al, Ba, Cd, Co, Cr, Cu, Mn, Ni, Th, Tl, and Zn). LIBS technology also showed the presence of a group of toxic and radioactive elements, which are as follows (Ac, Pu, Th, U) and (Al, Ba, Cd, Co, Cr, Cu, Hg, Mn, Ni, Pb, Tl, Zn). FTIR technique indicated the possibility of these items in of LNG project in Balhaf. Quantitative measurements of toxic metals have been carried out by using ICP, the results were compared with different international standards. By analyzing the results of the ICP for the samples, it is clear that the concentration of toxic elements is high. The LIBS technique showed the presence of the dangerous element the plutonium in all samples, with a higher intensity than all other elements.

#### REFERENCES

- [1] M. K. Hill, Understanding environmental pollution. Cambridge University Press, 2020.
- [2] J. J. Peirce, P. A. Vesilind, and R. Weiner, Environmental pollution and control. Butterworth-Heinemann, 1998.
- [3] I. L. Pepper, C. P. Gerba, and M. L. Brusseau, Environmental and pollution science. Elsevier, 2011.
- [4] C. Su, "A review on heavy metal contamination in the soil worldwide: Situation, impact and remediation techniques," Environ. Skept. Critics, vol. 3, no. 2, p. 24, 2014.
- [5] C. A. T. Force, "Fossil Fumes: A Public Health Analysis of Toxic Air Pollution From the Oil and Gas Industry (June 2016)."
- [6] L. M. McKenzie, R. Z. Witter, L. S. Newman, and J. L. Adgate, "Human health risk assessment of air emissions from development of unconventional



- natural gas resources," *Sci. Total Environ.*, vol. 424, pp. 79–87, 2012.
- [7] L. B. Paulik et al., "Impact of natural gas extraction on PAH levels in ambient air," *Environ. Sci. Technol.*, vol. 49, no. 8, pp. 5203–5210, 2015.
- [8] T. Colborn, C. Kwiatkowski, K. Schultz, and M. Bachran, "Natural gas operations from a public health perspective," *Hum. Ecol. risk Assess. An Int. J.*, vol. 17, no. 5, pp. 1039–1056, 2011.
- [9] F. Perera, "Pollution from fossil-fuel combustion is the leading environmental threat to global pediatric health and equity: Solutions exist," *Int. J. Environ. Res. Public Health*, vol. 15, no. 1, p. 16, 2018.
- [10] N. Fann et al., "Assessing human health PM2.5 and ozone impacts from US oil and natural gas sector emissions in 2025," *Environ. Sci. Technol.*, vol. 52, no. 15, pp. 8095–8103, 2018.
- [11] A. P. Pacsi, Y. Kimura, G. McGaughey, E. C. McDonald-Buller, and D. T. Allen, "Regional ozone impacts of increased natural gas use in the Texas power sector and development in the Eagle Ford shale," *Environ. Sci. Technol.*, vol. 49, no. 6, pp. 3966–3973, 2015.
- [12] C.-H. Chaîneau et al., "Results and lessons learned from a 3-year intensive coral communities monitoring during the construction of a LNG plant in Yemen," 2010.
- [13] "FACTBOX - Five facts about," vol. 15, p. 2010, 2010.
- [14] W. R. True, "Yemen LNG yields first production into glutted market," pp. 7–8, 2009.
- [15] W. L. Lindsay, *Chemical equilibria in soils*. John Wiley and Sons Ltd., 1979.
- [16] G. H. Ong, C. K. Yap, M. Mahmood, S. G. Tan, and S. Hamzah, "Barium levels in soils and *Centella asiatica*," *Trop. life Sci. Res.*, vol. 24, no. 1, p. 55, 2013.
- [17] A. M. Evans, W. L. Barrett, T. Bell, J. Milsom, C. J. Moon, and B. C. Scott, "Introduction to mineral exploration." Blackwell, London, 1993.
- [18] C. J. Moon, K. G. Whateley, and A. M. Evans, "Introduction to Mineral Exploration BLACKWELL PUBLISHING," New York, 2006.
- [19] H. E. Hawkes and J. S. Webb, *Geochemistry in mineral exploration*. Harper & Row, 1965.
- [20] J. E. Sansonetti and W. C. Martin, "Handbook of basic atomic spectroscopic data," *J. Phys. Chem. Ref. Data*, vol. 34, no. 4, pp. 1559–2259, 2005, doi: 10.1063/1.1800011.
- [21] W. H. O. WHO, "Mercury in Drinking-water, Background document for development of WHO Guidelines for Drinking-water Quality," *Who*, vol. WHO/SDE/WS, p. WHO/SDE/WSH/05.08/10, 2005, [Online]. Available: [http://www.who.int/water\\_sanitation\\_health/dwq/chemicals/mercuryfinal.pdf](http://www.who.int/water_sanitation_health/dwq/chemicals/mercuryfinal.pdf).
- [22] G. Gauglitz and D. S. Moore, *Handbook of spectroscopy*, vol. 1. Wiley-Vch Weinheim, 2014.
- [23] C. E. Fonrobert, "The political symbolism of the Eruv," *Jew. Soc. Stud.*, vol. 11, no. 3, pp. 9–35, 2005.
- [24] D. Lin-Vien, N. B. Colthup, W. G. Fateley, and J. G. Grasselli, *The handbook of infrared and Raman characteristic frequencies of organic molecules*. Elsevier, 1991.
- [25] Y. S. Mary et al., "Vibrational spectroscopic studies and computational study of 4-fluoro-N-(2'-hydroxy-4'-nitrophenyl) phenylacetamide," *J. Mol. Struct.*, vol. 994, no. 1–3, pp. 223–231, 2011.
- [26] A. Anantharaman, S. Ramalakshmi, and M. George, "Green synthesis of calcium oxide nanoparticles and its applications," *Int. J. Eng. Res. Appl*, vol. 6, no. 10, pp. 27–31, 2016.
- [27] B. H. Stuart, *Infrared spectroscopy: fundamentals and applications*. John Wiley & Sons, 2004.
- [28] L. J. Bellamy, "The Infrared Spectra of Complex Molecules" London Chapman Hall, 1980.